



- **FACT SHEET No. 15**

## **Temporomandibular Disorders**

### **Introduction**

Temporomandibular disorders (TMD) encompass a group of musculoskeletal and neuromuscular conditions that involve the temporomandibular joint (TMJ), the masticatory muscles, and all associated tissues. Pain associated with TMD can be clinically expressed as masticatory muscle pain (MMP) referred to as a myogenous temporomandibular disorder or TMJ pain (synovitis, capsulitis, osteoarthritis) referred to as an arthrogenous temporomandibular disorder. Chewing or other mandibular activity aggravates musculoskeletal pain. TMD pain can be (but is not necessarily) associated with biomechanical dysfunction of the temporomandibular joint (clicking or locking of the TMJ and limitation of jaw movement).

### **Epidemiology and Economics**

TMD-related facial pain has been reported in 9–13% of the general population (with a female: male ratio of 2:1), but only 4–7% seek treatment (female:male ratio of 4:1). Signs and symptoms peak among 20–40-year-olds. Incidence is approximately 3.9% per annum, and available data indicates that incidence among females is only slightly greater than among males and that it increases across age through the mid-forties, but is not known for older individuals. Progression to severe and/or chronic pain is associated with greater psychosocial distress, sleep disturbances, and comorbidities. TMD-related pain can affect daily activities, physical and psychosocial functioning, and quality of life.

Disabling TMD pain results in significant loss of work days and in other health-care-related costs.

### **Pathophysiology**

Many aspects of the etiology of TMD are unclear. In contrast to dental/occlusal causes, for which supporting evidence has been inadequate, there is definite support for a biopsychosocial and multifactorial background, illustrating the complex interaction between biological (e.g., hormonal) mechanisms, psychological states and traits, environmental conditions, and macro- and microtrauma.



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**

In MMP, experts propose a complex interaction between environmental, emotional, behavioral, and physical factors such as parafunctions including clenching or non forceful but frequent tooth contact during waking hours and or low level sustained muscle activity during sleep, (micro-) trauma, and release of inflammatory mediators and neuropeptides in muscles, which can sensitize the peripheral and central nervous systems. In conjunction with altered pain-regulating mechanisms (also influenced by female hormones), such factors may lead to localized or more generalized muscle pain, which is often associated with comorbidities. Recent articles have highlighted the cultural effects of persistent TMD pain on patient behavior, as well as genetic factors (COMT gene haplotypes).

TMJ arthralgia may result from trauma and from intrinsic and extrinsic overloading of the TMJ that may overcome the adaptive capacity of the joint tissues; this seem to be more critical for osteoarthritis of TMJ. Alternatively, the adaptive capacity of the TMJ may be reduced by intrinsic factors such as reduced blood supply and inadequate nutrition. Genetics and gender have also been implicated in the pathophysiology of osteoarthritis. The production of free radicals, proinflammatory and nociceptive neuropeptides, enzymes, bone morphogenetic proteins, and growth factors will lead to inflammation, pain, and progressive tissue changes.

### **Clinical Features**

Myogenous temporomandibular disorder or MMP is a regional, dull, aching pain, most prominent in the jaw-closing muscles, which can occur at rest and may be aggravated during mandibular function. Pain may be more pronounced in the morning or evening and ranges from mild to severe in intensity. Reported associated symptoms are limitation of movement, headache, fullness of the ear, and neck pain, but cause-and-effect relationships have not yet been established. The regional disorder of MMP should be distinguished from MMP that occurs as part of a generalized muscle pain disorder such as fibromyalgia.

Arthrogenous temporomandibular disorder or TMJ arthralgia is a more localized and sharp pain of moderate to severe intensity, localized to the TMJ and surrounding tissues and radiating mostly to the ear region. The pain is aggravated during loading and functioning of the joint. In addition, there may be limited movement and function. TMJ pain may be associated with a displaced or dysfunctional articular disk causing joint locking, which may be an additional cause of limited movement. TMJ osteoarthritis may be part of generalized arthritis and is accompanied by crepitation.

If chronic pain develops, both myogenous and arthrogenous temporomandibular disorders may be accompanied by central sensitization and psychological problems such as depression, somatization, and anxiety.

### **Diagnostic Criteria**

The Guidelines of the American Academy of Orofacial Pain (2013) and the Diagnostic Criteria for Temporomandibular Disorders (DC-TMD), suggest the following criteria:

MMP: A complaint of muscle pain in the jaw, in the temple, in the ear, or in front of the ear that is affected by jaw movement, function, or parafunction. Replication of this familiar pain occurs with



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**

provocation testing of the masticatory muscles (i.e., palpation of the temporalis or masseter muscle(s); OR with maximum unassisted or assisted opening. Limitation of mandibular movement(s) secondary to pain may be present.

TMJ arthralgia: A complaint of joint pain that is affected by jaw movement, function, or parafunction. Replication of this familiar pain occurs with provocation testing of the TMJ (i.e., palpation of the lateral pole or around the lateral pole) OR with maximum unassisted or assisted opening, right or left lateral movements, or protrusive movements.

Psychosocial factors are rated by means of a pain drawing for pain locations and comorbidities, the Graded Chronic Pain Scale (GCPS) for pain intensity and physical function, the Jaw Function Limitation Scale (JFLS) short-form for limited function, the Patient History Questionnaire-4 (PHQ-4) for depression and anxiety, and the Oral Behavior Checklist for parafunction.

The DC/TMD Criteria are specific for various diagnostic paradigms beyond the scope of this fact sheet. More information may be obtained at: <http://www.rdc-tmdinternational.org/TMDAssessmentDiagnosis/DCTMD.aspx>

### **Diagnosis and Treatment**

The gold standard for a pain-related TMD diagnosis is the combination of history and clinical examination. Except for imaging, technical examinations (e.g., electromyography or occlusal analysis) are not warranted. The symptoms are usually self-limiting, with a benign natural course.

Management aims at providing the optimal circumstances for healing and adaptation to take place.

Noninvasive, reversible therapies that fit in the biopsychosocial approach include:

- Education of the patient, active self-care, follow-up, medical monitoring
- Physical therapy, behavioral self-regulation programs
- Intraoral occlusal appliances
- Medication (analgesics, nonsteroidal anti-inflammatory drugs)

In patients with chronic TMD, these therapies must be accompanied by:

- Psychological support, e.g., cognitive-behavioral therapy, relaxation therapy
- Low-dose tricyclic antidepressants

For patients with persistent TMJ arthralgia, arthrocentesis might be considered, but TMJ surgery is rarely, if ever, indicated in the scope of TMD pain treatment.

### **REFERENCES**

1. Benoliel R, Sharav Y. Masticatory myofascial pain, and tension-type and chronic daily headache. In: Sharav Y, Benoliel R, editors. Orofacial pain and headache. Edinburgh: Elsevier; 2008. p. 109-28.
2. Benoliel R, Svensson P, Heir GM, Sirois D, Zakrzewska J, Oke-Nwosu J, Torres SR, Greenberg MS, Klasser GD, Katz J, Eliav E. Persistent orofacial muscle pain. Oral Dis 2011;17(Suppl 1):23-41.



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**

3. Chen CY, Palla S, Erni S, Sieber M, Gallo LM. Nonfunctional tooth contact in healthy controls and patients with myogenous facial pain. *J Orofac Pain*. 2007 Summer;21(3):185-93.
4. De Boever JA, Nilner M, Orthlieb JD, Steenks MH; Educational Committee of the European Academy of Craniomandibular Disorders. Recommendations by the EACD for examination, diagnosis, and management of patients with temporomandibular disorders and orofacial pain by the general dental practitioner. *J Orofac Pain* 2008;22:268–78.
5. de Leeuw R, Klasser G. Orofacial pain. Guidelines for assessment, diagnosis and management, 5th ed. The American Academy of Orofacial Pain. Quintessence; 2013.
6. Dubrovsky B, Raphael KG, Lavigne GJ, Janal MN, Sirois DA, Wigren PE, Nemelivsky LV, Klausner JJ, Krieger AC. Polysomnographic investigation of sleep and respiratory parameters in women with temporomandibular pain disorders. *J Clin Sleep Med*. 2014 Feb 15;10(2):195-201. doi: 10.5664/jcsm.3452.
7. Funato M, Ono Y, Baba K, Kudo Y. Evaluation of the non-functional tooth contact in patients with temporomandibular disorders by using newly developed electronic system. *J Oral Rehabil*. 2014 Mar;41(3):170-6.
8. Greene CS. Managing the care of patients with temporomandibular disorders: a new guideline for care. *J Am Dent Assoc* 2010;141:1086–8.
9. List T, Axelsson S. Management of TMD: evidence from systematic reviews and meta-analyses. *J Oral Rehabil* 2010;37:430–51.
10. Manfredini D, Guarda-Nardini L, Winocur E, Piccotti F, Ahlberg J, Lobbezoo F. Research diagnostic criteria for temporomandibular disorders: a systematic review of axis I epidemiologic findings. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:453–62.
11. Manfredini D, Winocur E, Guarda-Nardini L, Lobbezoo F. Epidemiology of bruxism in adults: a systematic review of the literature. *J Orofac Pain* 2013;27:99–110.
12. Michelotti A, Liguori R, Toriello M, D'Antò V, Vitale D, Castaldo G, Sacchetti L. Catechol-O-methyltransferase (COMT) gene polymorphisms as risk factor in temporomandibular disorders patients from southern Italy. *Clin J Pain* 2013; Epub Feb 26.
13. Raphael KG, Janal MN, Sirois DA, Dubrovsky B, Wigren PE, Klausner JJ, Krieger AC, Lavigne GJ. Masticatory muscle sleep background electromyographic activity is elevated in myofascial temporomandibular disorder patients. *J Oral Rehabil*. 2013 Dec;40(12):883-91.
14. Raphael KG, Sirois DA, Janal MN, Wigren PE, Dubrovsky B, Nemelivsky LV, Klausner JJ, Krieger AC, Lavigne GJ. Sleep bruxism and myofascial temporomandibular disorders: a laboratory-based polysomnographic investigation. *J Am Dent Assoc*. 2012 Nov;143(11):1223-31.
15. Sanders AE, Essick GK, Fillingim R, Knott C, Ohrbach R, Greenspan JD, Diatchenko L, Maixner W, Dubner R, Bair E, Miller VE, Slade GD. Sleep apnea symptoms and risk of temporomandibular disorder: OPPERA cohort. *J Dent Res*. 2013 Jul;92(7 Suppl):70S-7S.
16. Schiffman E, Ohrbach R, Truelove E, Look J, Anderson G, Goulet JP, List T, Svensson P, Gonzalez Y, Lobbezoo F, Michelotti A, Brooks SL, Ceusters W, Drangsholt M, Ettlin D, Gaul C, Goldberg LJ, Haythornthwaite JA, Hollender L, Jensen R, John MT, De Laat A, de Leeuw R, Maixner W, van der Meulen M, Murray GM, Nixdorf DR, Palla S, Petersson A, Pionchon P, Smith B, Visscher CM, Zakrzewska J, Dworkin SF; International RDC/TMD Consortium Network, International association for Dental Research; Orofacial Pain Special Interest Group, International Association for the Study of Pain; Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: recommendations of the International RDC/TMD Consortium Network\* and Orofacial Pain Special Interest Group; *J Oral Facial Pain Headache*. 2014 Winter;28(1):6-27. doi: 10.11607/jop.1151.
17. Schmitter A, Kares-Vrincianu A, Kares H, Bermejo JL, Schindler HJ. Sleep-associated aspects of myofascial pain in the orofacial area among Temporomandibular Disorder patients and controls. *Sleep Med*. 2015 Sep;16(9):1056-61
18. Slade GD, Bair E, Greenspan JD, Dubner R, Fillingim RB, Diatchenko L, Maixner W, Knott C, Ohrbach R. Signs and symptoms of first-onset TMD and socio-demographic predictors of its development: the OPPERA prospective cohort study. *Journal of Pain* 2013. 14(Supple2):T20-T32.
19. Wang XD, Zhang JN, Gan YH, Zhou Y Current understanding of pathogenesis and treatment of TMJ osteoarthritis. *J Dent Res*. . 2015. 94(5):666–673.

### **About the International Association for the Study of Pain®**

IASP is the leading professional forum for science, practice, and education in the field of pain. [Membership is open to all professionals](#) involved in research, diagnosis, or treatment of pain. IASP has more than 7,000 members in 133 countries, 90 national chapters, and 20 Special Interest Groups.

**As part of the Global Year Against Musculoskeletal Pain, IASP offers a series of Fact Sheets that cover specific topics related to postsurgical pain. These documents have been translated into multiple languages and are available for free download. Visit [www.iasp-pain.org/globalyear](http://www.iasp-pain.org/globalyear) for more information.**



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**